Validation data for RAW-Blue™ Cells

https://www.invivogen.com/raw-blue

For research use only

Version 23B16-AK

RAW-BlueTM cells are designed to monitor the NF- κ B and AP-1 responses upon PRR (pattern recognition receptor) stimulation. These cells stably express an NF- κ B/AP-1-inducible secreted embryonic alkaline phosphatase (*SEAP*) reporter gene. Levels of SEAP activities are readily assessable in the supernatant using QUANTI-BlueTM Solution, a SEAP detection medium. RAW-BlueTM cells are derive from the murine RAW 264.7 macrophage cell line, which has been reported to express many pattern recognition receptors (PRRs), as verified using RT-PCR (Figure 1) and by monitoring the NF- κ B-response upon stimulation using various PRR-ligands (Figure 2).

Validation of PRR expression




Figure 1: Expression profile of various PRRs in RAW-Blue[™] cells. RT-PCR results show endogenous expression of Toll-like receptors TLRs (1-9), RIG-I, MDA-5, NOD1, NOD2 and Dectin-1 in RAW-Blue[™] cells.




Figure 2: NF-κB responses in RAW-Blue™ cells to various PRR ligands. Cells were incubated with TLR, Dectin and NOD agonists: TLR2 (HKLM, 1x108 cells/ml), TLR1/2 (Pam3CSK4, 1 μg/ml), TLR2/6 (FSL-1, 1 μg/ml), TLR3 (Poly(I:C) HMW, 100 μg/ml), TLR4 (LPS-EK, 10 μg/ml), TLR5 (FLA-ST Ultrapure (UP), 10 μg ml), TLR7 (CL075, 10 μg/ml), TLR9 (ODN1826, 10 μg/ ml), NOD1 (Tri-DAP, 10 μg/ml), NOD2 (MDP, 10 μg/ml) and Dectin-1 (Zymosan, 100 μg/ml). After 24h incubation, TLR and NOD stimulation was assessed by measuring the levels of SEAP using QUANTI-Blue™. Data are shown as optical density (OD) at 630 nm (mean ± SEM).

TECHNICAL SUPPORT InvivoGen USA (Toll-Free): 888-457-5873

InvivoGen USA (International): +1 (858) 457-5873 InvivoGen Europe: +33 (0) 5-62-71-69-39 InvivoGen Asia: +852 3-622-34-80

E-mail: info@invivogen.com

